

YEAR 9 PHYSICS, AQA GCSE, UNIT 2, PART 1

ENERGY STORES AND TRANSFERS

PART 1 OF 3

LESSONS 1–3: ENERGY STORES AND SYSTEMS, THE KINETIC STORE, THE GRAVITATIONAL POTENTIAL STORE

3

LESSONS


1

PRACTICAL

3

NEW EQUATIONS

ANSWER EDITION FOR TEACHERS AND SELF-MARKING



NAME

TEACHER

CLASS

TARGET GRADE

HOW THIS BOOKLET WORKS

Energy is never made and never destroyed: it only moves from one store to another. This booklet shows you how to follow it, and how to calculate it. Keep the booklet: it is your classwork book *and* your revision guide.



DO NOW

Quick questions from earlier lessons. Remembering again is what makes it stick.



LEARN IT

The explanation, diagrams and worked examples. Read this again when you revise.



TRY IT

Practice that builds up: first with support, then on your own.



PRACTICAL

Record your measurements and work out results from them.



EXAM CORNER

AQA-style questions with a tip on how the marks are given.



REVISE IT

Summary, cover-and-answer questions and an “I can” check to colour in.



Two kinds of question to know. When a sentence has a gap, the choices are printed straight after it, like this: Energy is measured in joules (joules / newtons / watts). When you see a box of choices, **draw a ring around** the correct one, as AQA exam papers ask. The unit of energy is the watt the joule the newton .

THE ROUTE THROUGH THIS PART

#	LESSON	WEEK
01	Energy stores and systems The energy stores, four ways to transfer energy, systems	1
02	The kinetic store $E_k = \frac{1}{2} m v^2$; grams to kilograms; double the speed, four times the energy	1
03	The gravitational potential store What g means, $E_p = m g h$, height gained, finding h or m	2

This booklet is Part 1 of 3: Lessons 1–3. The **equation toolkit** on the last page has every equation you have met so far, with formula triangles to help you rearrange.

ENERGY STORES AND SYSTEMS

AQA Physics 4.1.1.1; Combined Science 6.1.1.1

BIG QUESTION A skateboarder rolls down one side of a ramp and up the other, but never quite gets back to the starting height. Where did the energy go?

BY THE END YOU CAN

- name the energy stores and give an example of each
- name the four ways energy is transferred
- explain what a system is
- describe the energy transfers when something changes

energy store

transfer

system

closed system

joule (J)

YOU WILL NEED

This booklet, a pencil and a ruler.



REMEMBER

WHAT DO YOU ALREADY THINK?

No wrong answers: this is your starting point.

1

Name something that has energy. How can you tell?

For example, a moving car (it is moving) or a hot drink (it is hot). Any sensible answer.

2

Energy is measured in joules (joules / newtons / metres).

joules (J)

3

Rub your hands together. They get warmer (warmer / colder). Why?

Warmer: rubbing (friction) transfers energy to the thermal store of your hands.



Energy is never made and never used up. It is **stored**, and it is **transferred** (moved) from one store to another. Energy is measured in **joules (J)**.

■ THE FIVE MAIN ENERGY STORES

STORE	ENERGY IS STORED BECAUSE...	EXAMPLE
Kinetic	the object is moving	a rolling football
Gravitational potential	the object is raised up	a book on a high shelf
Elastic potential	the object is stretched or squashed	a stretched spring, a drawn bow
Thermal	the object is hot	a cup of tea
Chemical	of the substances it is made of	food, fuel, a battery

You may also meet the **magnetic** store (two magnets held apart), the **electrostatic** store (a charged balloon) and the **nuclear** store (uranium fuel), but GCSE questions rarely use them.

■ FOUR WAYS TO TRANSFER ENERGY

TRANSFERRED...	WHAT HAPPENS	EXAMPLE
mechanically	a force moves something	pushing a trolley
electrically	a current flows	a battery lighting a lamp
by heating	a hotter object warms a colder one	a hob heating a pan
by waves	a wave carries it: light, infrared (both called radiation) or sound	the Sun warming your face; a loudspeaker



Stores hold energy. Transfers move it. Light, sound and electricity are **not** stores: they are ways energy travels from one store to another.

■ SYSTEMS

A **system** is the object, or group of objects, that we choose to study. In a **closed system**, no energy gets in or out, so the **total** energy stays the same. The energy only moves between stores.

■ DESCRIBE ANY CHANGE IN THREE STEPS

- 1 Which store **empties**?
- 2 Which store **fills**?
- 3 **How** is the energy transferred: mechanically, electrically, by heating or by waves?

EXAMPLE: A CAR BRAKING


The kinetic store of the car empties. The thermal store of the brakes fills. Friction is a force, so the energy is transferred mechanically. **Write:** "Energy is transferred mechanically from the kinetic store of the car to the thermal store of the brakes."



PRACTISE

TRY IT

- 1 Draw a line from each store to its example.



- 2 Draw a ring around the correct answer in each box.

Light is an energy store a way energy is transferred

A battery stores energy in its chemical store kinetic store thermal store


At the top of its flight, a ball has most energy in its

kinetic store gravitational potential store


- 3 The pictures tell a story. For each picture, write the store that **empties** and the store that **fills**. Choose from the words under each gap.




Eating breakfast



Cycling up the hill



Freewheeling down



Braking at the bottom

PICTURE	STORE THAT EMPTIES	STORE THAT FILLS
A Eating	chemical (in the food)	chemical (in the body)
B Cycling up	chemical / kinetic chemical (body)	gravitational potential / elastic potential gravitational potential (and thermal)
C Freewheeling down	gravitational potential / chemical gravitational potential	kinetic / nuclear kinetic
D Braking	kinetic / thermal kinetic	thermal / chemical thermal (brakes and surroundings)

- 4 **SUPPORTED** Complete each sentence. The choices are after each gap.

A ball is thrown upwards. Energy is transferred from the kinetic (kinetic / chemical) store to the gravitational potential (gravitational potential / thermal) store.

A fireplace warms a person. Energy is transferred by infrared waves from the chemical (chemical / kinetic) store of the fuel to the thermal (thermal / kinetic) store of the person.

A child lands on a trampoline. Energy is transferred mechanically from the child's kinetic (kinetic / chemical) store to the trampoline's elastic potential (elastic potential / thermal) store.

- 5 **ON YOUR OWN** A bungee jumper falls, and the cord stretches until the jumper stops at the bottom. Describe the energy transfers. Use the three steps.

As the jumper falls, energy is transferred from the gravitational potential store of the jumper to the kinetic store. As the cord stretches, energy is transferred from the kinetic store to the elastic potential store of the cord.

6 Explain what is wrong with what Sam says.

Sam

When the car stopped, its energy was used up.

Energy cannot be used up (destroyed). The energy in the car's kinetic store was transferred to the thermal store of the brakes and the surroundings.



AQA-STYLE

EXAM CORNER

Write in full sentences. Show every step of a calculation.

Get the marks: Name both stores and say which one empties and which one fills.

7 A student throws a ball straight up. Complete the sentence. [2 marks]

As the ball rises, energy is transferred from the kinetic (kinetic / chemical / thermal) store of the ball to its gravitational potential (elastic potential / gravitational potential / nuclear) store.

8 What is the unit of energy? Tick **one** box. [1 mark]

☒

joule

☐

newton

☐

watt

☐

kilogram

9 A child lands on a trampoline. Describe the energy transfer as the mat and springs stretch and the child slows down. [2 marks]

Energy is transferred mechanically from the kinetic store of the child (1) to the elastic potential store of the trampoline (1).



SUMMARY

- Energy is measured in **joules (J)**. It is never made or destroyed.
- Main stores: kinetic, gravitational potential, elastic potential, thermal, chemical.
- Transfers: mechanically, electrically, by heating, by waves.
- In a closed system, the total energy stays the same.

COVER THE PAGE AND ANSWER

1 What is the unit of energy?
.....

2 Name four energy stores.
.....

3 Is light a store or a way of transferring energy?
.....

4 What is a closed system?
.....

I CAN...

name the energy stores



name the four ways energy is transferred



explain what a system is



describe energy transfers using "from ... to ..."



THE KINETIC STORE

AQA Physics 4.1.1.2; Combined Science 6.1.1.2

BIG QUESTION A car travels twice as fast. Does it have twice as much energy in its kinetic store, or more?

BY THE END YOU CAN

- say what the kinetic store depends on
- calculate kinetic energy with $E_k = \frac{1}{2} m v^2$
- convert grams to kilograms
- explain why doubling the speed gives four times the energy

kinetic store

mass (kg)

speed (m/s)

squared

YOU WILL NEED

Calculator. Practical: trolley, 1.00 m track, stopwatch, balance.



REMEMBER

DO NOW

From earlier lessons. No notes!

1

Which store does a moving object have energy in?

The kinetic store.

2

A car brakes. Which store fills?

The thermal store (of the
brakes and the
surroundings).

3

Light is

a way energy is
transferred

(a store / a way energy is
transferred).

A way energy is

transferred (by waves).



EXPLAIN

ENERGY IN A MOVING OBJECT

Anything that moves has energy in its **kinetic store**. The more **mass** it has, and the faster it moves, the more energy it has.

$$E_k = \frac{1}{2} m v^2$$

kinetic energy = 0.5 × mass × (speed)²

E_k in joules (J), m in kilograms (kg), v in metres per second (m/s)

Only the speed is squared. In this unit you use the equation to calculate the energy.



Square the speed first: $v^2 = v \times v$. Then multiply by the mass and by 0.5.

WORKED EXAMPLE

A 1200 kg car moves at 15 m/s. Calculate the energy in its kinetic store.

KNOW $m = 1200 \text{ kg}, v = 15 \text{ m/s}$

EQUATION $E_k = \frac{1}{2} m v^2$

SUBSTITUTE $E_k = 0.5 \times 1200 \times 15^2$

ANSWER $E_k = 135\,000 \text{ J}$

YOUR TURN

A 60 kg runner moves at 5.0 m/s. Calculate the energy in their kinetic store.

KNOW $m = 60 \text{ kg}, v = 5.0 \text{ m/s}$

EQUATION $E_k = \frac{1}{2} m v^2$

SUBSTITUTE $E_k = 0.5 \times 60 \times 5.0^2$

ANSWER $E_k = 750 \text{ J}$

MASS IN GRAMS? CONVERT FIRST

The mass must be in kilograms. Divide grams by 1000: $450 \text{ g} = 0.450 \text{ kg}$. A 450 g football kicked at 20 m/s has $E_k = 0.5 \times 0.450 \times 20^2 = 90 \text{ J}$.

DOUBLE THE SPEED, FOUR TIMES THE ENERGY

SPEED OF A 1200 KG CAR	v^2	E_k
15 m/s	225	135 000 J
30 m/s (2 × faster)	900	540 000 J (4 × more)
45 m/s (3 × faster)	2025	1 215 000 J (9 × more)

Twice the speed gives $2^2 = 4$ times the energy. Three times the speed gives $3^2 = 9$ times. This is why a crash at high speed does so much more damage, and why the brakes must transfer so much more energy to stop the car. Doubling the **mass** only doubles the energy, because the mass is not squared.



PRACTISE
TRY IT

- 1 Draw a ring around the correct answer in each box.

Doubling the mass makes the kinetic energy

stay the same **2 times bigger** 4 times bigger

Doubling the speed makes the kinetic energy

stay the same 2 times bigger **4 times bigger**

- 2 **SUPPORTED** A 2.0 kg ball rolls at 3.0 m/s. Calculate the energy in its kinetic store.

EQUATION	$E_k = \frac{1}{2} m v^2$	
SUBSTITUTE	$E_k = 0.5 \times 2.0 \times 3.0^2$	
ANSWER	9.0 J	unit:

- 3 **SUPPORTED** A 450 g football is kicked at 20 m/s. Calculate the energy in its kinetic store.
Convert the mass first!

CONVERT	$m = 450 \div 1000 = 0.450 \text{ kg}$	
EQUATION	$E_k = \frac{1}{2} m v^2$	
SUBSTITUTE	$E_k = 0.5 \times 0.450 \times 20^2$	
ANSWER	90 J	unit:

- 4 Complete the table. Show your working underneath.

Object	Mass	Speed	E_k / J
Lorry	30 000 kg	2.0 m/s	60 000
Cyclist and bike	80 kg	6.0 m/s	1440
Tennis ball	58 g	50 m/s	72.5
Bullet	10 g	400 m/s	800

Working area for calculations.

5

ON YOUR OWN

A 0.16 kg cricket ball is bowled at 40 m/s. Calculate the energy in its kinetic store.

$$E_k = 0.5 \times 0.16 \times 40^2 = 128 \text{ J}$$

6

CHALLENGE

A car speeds up from 10 m/s to 30 m/s. How many times bigger is the energy in its kinetic store? Explain without calculating the energy.

9 times bigger. The speed is 3 times bigger, and the speed is squared: $3^2 = 9$.

7

Explain what is wrong with what Ava says.

Ava

At 4 m/s my bike has twice as much kinetic energy as at 2 m/s.

The speed is squared, so doubling the speed gives $2^2 = 4$ times as much kinetic energy, not twice as much.



MEASURE THE ENERGY OF A MOVING TROLLEY

Work in a group of three: one pushes, one times, one records.

- 1 Measure the trolley's mass on the balance in grams. Convert it to kilograms.
- 2 Mark a 1.00 m section of the track with tape.
- 3 Give the trolley a gentle push so it rolls through the section. Time how long it takes to cross the 1.00 m.
- 4 Repeat twice more with the same push, and find the mean time.
- 5 Calculate the speed, then the kinetic energy.
- 8 Record your results.

Mass / g	Mass / kg	Time 1 / s	Time 2 / s	Time 3 / s	Mean time / s
500	0.500	1.62	1.55	1.59	1.59

- 9 Calculate the speed: $\text{speed} = \frac{\text{distance}}{\text{time}}$.

EQUATION	$v = \frac{s}{t}$	
SUBSTITUTE	$v = \frac{1.00}{1.59}$	
ANSWER	0.63 m/s (sample)	unit:

- 10 Calculate the energy in the trolley's kinetic store.

EQUATION	$E_k = \frac{1}{2} m v^2$	
SUBSTITUTE	$E_k = 0.5 \times 0.500 \times 0.63^2$	
ANSWER	0.10 J (sample)	unit:

- 11 Which measurement has the biggest effect on your answer?

the mass the time

Explain why.

The time: the speed is worked out from it, and the speed is squared, so any error in the time is made bigger.



Get the marks: Write the equation, then substitute, then give the answer with its unit.

12

A cyclist and bike have a total mass of 70 kg and move at 8.0 m/s.
Calculate the energy in their kinetic store.

[2 marks]

EQUATION	$E_k = \frac{1}{2} m v^2$	
SUBSTITUTE	$E_k = 0.5 \times 70 \times 8.0^2$ (1)	
ANSWER	2240 J (1)	unit:

13

Car A and car B have the same mass. Car B moves twice as fast as car A. Complete the sentence.

[2 marks]

Car B has 4 (2 / 4 / 8) times as much energy in its kinetic store as car A, because in the equation the speed is halved squared doubled.

**SUMMARY**

- $E_k = \frac{1}{2} m v^2$: mass in kg, speed in m/s, energy in J.
- Convert grams to kilograms: divide by 1000.
- Only the speed is squared: 2 × the speed gives 4 × the energy.
- 2 × the mass gives 2 × the energy.

COVER THE PAGE AND ANSWER

1 What two things does the kinetic store depend on?

2 Write the equation for kinetic energy.

3 Convert 250 g to kilograms.

4 The speed triples. What happens to the kinetic energy?

I CAN...

calculate kinetic energy with $E_k = \frac{1}{2} m v^2$



convert grams to kilograms



explain why doubling the speed gives four times the energy



THE GRAVITATIONAL POTENTIAL STORE

AQA Physics 4.1.1.2; Combined Science 6.1.1.2

BIG QUESTION Carrying a bag up the stairs is hard work. Where does that energy go, and can you get it back?

BY THE END YOU CAN

- explain what $g = 9.8 \text{ N/kg}$ means
- calculate energy with $E_p = mgh$
- use the height gained
- rearrange to find a height or a mass

gravitational potential store

gravitational field strength

newton (N)

height gained

ELECTRIC MOUNTAIN

At Dinorwig in Wales, water is pumped up into a lake high on a mountain. When the country suddenly needs electricity, the water rushes down through turbines. The energy waits in the water's gravitational potential store until it is needed.



REMEMBER

DO NOW

From earlier lessons. No notes!

1

In $E_k = \frac{1}{2}mv^2$, what does v stand for? Give its unit.

Speed, in metres per

second (m/s).

2

Convert 600 g to kilograms.

0.600 kg

3

A 2.0 kg ball moves at 4.0 m/s. Calculate its kinetic energy.

$$E_k = 0.5 \times 2.0 \times 4.0^2$$

$$E_k = 16 \text{ J}$$




EXPLAIN

ENERGY FROM HEIGHT

Lift an object up and you transfer energy to its **gravitational potential store**. Let it fall and the energy is transferred to its kinetic store. The heavier the object and the higher you lift it, the more energy it stores.

WHAT g MEANS

! On Earth, gravity pulls every kilogram down with a force of **9.8 newtons**. We write this as $g = 9.8 \text{ N/kg}$ (newtons per kilogram). It is called the **gravitational field strength**. On the Moon, $g = 1.6 \text{ N/kg}$, so the same lift stores less energy there.




$$E_p = m g h$$

gravitational potential energy = mass $\times$
gravitational field strength $\times$ height

E_p in J, m in kg, g in N/kg, h in m

Rearranged: $h = \frac{E_p}{m g}$ and $m = \frac{E_p}{g h}$



WORKED EXAMPLE

A 2.0 kg book is lifted onto a shelf 1.5 m high. Calculate the energy transferred to its gravitational potential store.

KNOW $m = 2.0 \text{ kg}$, $g = 9.8 \text{ N/kg}$, $h = 1.5 \text{ m}$

EQUATION $E_p = m g h$

SUBSTITUTE $E_p = 2.0 \times 9.8 \times 1.5$

ANSWER $E_p = 29.4 \text{ J}$

YOUR TURN

A 5.0 kg bag is lifted 2.0 m. Calculate the energy transferred to its gravitational potential store.

KNOW $m = 5.0 \text{ kg}$, $g = 9.8 \text{ N/kg}$, $h = 2.0 \text{ m}$


EQUATION $E_p = m g h$

SUBSTITUTE $E_p = 5.0 \times 9.8 \times 2.0$

ANSWER $E_p = 98 \text{ J}$

USE THE HEIGHT GAINED

Only the **change** in height matters. A hiker climbs from a car park 120 m above sea level to a summit 420 m above sea level. The height gained is $h = 420 - 120 = 300 \text{ m}$. A 55 kg hiker gains $E_p = 55 \times 9.8 \times 300 = 161\,700 \text{ J}$.



■ FINDING THE HEIGHT OR THE MASS

WORKED EXAMPLE

A 60 kg climber gains 17 640 J in their gravitational potential store. How high did they climb?

KNOW $E_p = 17\,640\text{ J}$, $m = 60\text{ kg}$, $g = 9.8\text{ N/kg}$

REARRANGE $h = \frac{E_p}{m g}$

SUBSTITUTE $h = \frac{17\,640}{60 \times 9.8}$

ANSWER $h = 30\text{ m}$

YOUR TURN

A crate is lifted 2.0 m and gains 392 J. Calculate its mass.

KNOW $E_p = 392\text{ J}$, $g = 9.8\text{ N/kg}$, $h = 2.0\text{ m}$

REARRANGE $m = \frac{E_p}{g h}$

SUBSTITUTE $m = \frac{392}{9.8 \times 2.0}$

ANSWER $m = 20\text{ kg}$

Get the marks: Work out the bottom of the fraction first ($60 \times 9.8 = 588$), then divide. Or put brackets round it on your calculator.



PRACTISE

TRY IT

- 1 Draw a ring around the correct answer in each box.

On the Moon, lifting a box 1 m stores

more energy than on Earth **less energy than on Earth** the same energy as on Earth

Lifting a box twice as high makes the energy stored

half as big **twice as big** four times as big

- 2 **SUPPORTED** A 0.50 kg ball is lifted 3.0 m. Calculate the energy transferred to its gravitational potential store.

EQUATION	$E_p = m g h$	
SUBSTITUTE	$E_p = 0.50 \times 9.8 \times 3.0$	
ANSWER	14.7 J	unit:

3

SUPPORTED A 600 g book is lifted 1.2 m onto a shelf. Calculate the energy transferred.
Convert the mass first!

CONVERT	$m = 600 \div 1000 = 0.600 \text{ kg}$	
EQUATION	$E_p = m g h$	
SUBSTITUTE	$E_p = 0.600 \times 9.8 \times 1.2$	
ANSWER	7.1 J	unit:

4

Complete the table. Use $g = 9.8 \text{ N/kg}$. Show your working underneath.

Object	Mass / kg	Height gained / m	E_p / J
Child at the top of a slide	30	2.5	735
Rock on a cliff	4.0	5.0	196
Crate on a lorry	30	3.0	882
Lift with passengers	800	12	94 080

5

CHALLENGE A 50 kg student gains 4.9 kJ in their gravitational potential store climbing stairs. How high did they climb? Convert kJ to J first.

$$E_p = 4900 \text{ J}; h = \frac{E_p}{m g} = \frac{4900}{50 \times 9.8} = 10 \text{ m}$$



Choose three objects. Measure the mass of each and the height of the shelf or bench you lift it onto. Calculate the energy transferred to each gravitational potential store.

- 6 Record and calculate.

Object	Mass / kg	Height / m	E_p / J
textbook	1.2	0.90	10.6
water bottle	0.50	0.90	4.4
school bag	4.0	0.90	35.3

- 7 How high would you need to lift your lightest object so that it stores the same energy as your heaviest object?

$$\text{Sample: } h = \frac{E_p}{m g} = \frac{35.3}{0.50 \times 9.8} = 7.2 \text{ m}$$



Get the marks: g is always given in the question. AQA uses 9.8 N/kg.

- 8 A crane lifts a 250 kg load through a height of 12 m. Calculate the energy transferred to the load's gravitational potential store. Gravitational field strength = 9.8 N/kg **[2 marks]**

EQUATION	$E_p = m g h$	
SUBSTITUTE	$E_p = 250 \times 9.8 \times 12$ (1)	
ANSWER	29 400 J (1)	unit:

- 9 A 0.40 kg ball gains 7.84 J in its gravitational potential store when it is thrown upwards. Calculate how high it rises. Gravitational field strength = 9.8 N/kg **[3 marks]**

EQUATION	$E_p = m g h$	
REARRANGE	$h = \frac{E_p}{m g}$ (1)	
SUBSTITUTE	$h = \frac{7.84}{0.40 \times 9.8}$ (1)	
ANSWER	2.0 m (1)	unit:



SUMMARY

- $g = 9.8$ N/kg: gravity pulls each kilogram down with 9.8 N.
- $E_p = m g h$, with the height gained in metres.
- Rearranged: $h = \frac{E_p}{m g}$ and $m = \frac{E_p}{g h}$.
- Twice the height or twice the mass gives twice the energy.

COVER THE PAGE AND ANSWER

1 What does $g = 9.8$ N/kg mean?

2 Write the equation for gravitational potential energy.

3 A 5.0 kg bag is lifted 2.0 m. How much energy does it gain?

4 Rearrange $E_p = m g h$ to find h .

I CAN...

explain what $g = 9.8$ N/kg means



calculate $E_p = m g h$ using the height gained



rearrange $E_p = m g h$ to find a height or a mass



KEY WORDS

Learn the meaning and the spelling. In exams, precise words earn marks: say “transferred to the thermal store of the surroundings”, never “lost”.

Energy store [L1]: Where energy is held. The main stores are kinetic, gravitational potential, elastic potential, thermal and chemical

Joule (J) [L1]: The unit of energy

System [L1]: The object or group of objects that we choose to study

Closed system [L1]: A system that no energy enters or leaves; its total energy stays the same

Kinetic store [L2]: Energy an object has because it is moving

Gravitational potential store [L3]: Energy an object has because it is raised in a gravitational field

Gravitational field strength (g) [L3]: The force of gravity on each kilogram: 9.8 N/kg on Earth

PROGRESS TRACKER

Colour one circle for each statement: **red** = not yet, **amber** = getting there, **green** = I can do this without help. Revisit your reds first.

I CAN...	R / A / G
01: Energy stores and systems	
name the energy stores	☐ ☐ ☐
name the four ways energy is transferred	☐ ☐ ☐
explain what a system is	☐ ☐ ☐
describe energy transfers using "from ... to ..."	☐ ☐ ☐
02: The kinetic store	
calculate kinetic energy with $E_k = \frac{1}{2} m v^2$	☐ ☐ ☐
convert grams to kilograms	☐ ☐ ☐
explain why doubling the speed gives four times the energy	☐ ☐ ☐
03: The gravitational potential store	
explain what $g = 9.8$ N/kg means	☐ ☐ ☐
calculate $E_p = m g h$ using the height gained	☐ ☐ ☐
rearrange $E_p = m g h$ to find a height or a mass	☐ ☐ ☐

QUICK ANSWERS

Answers to the “Do now” and “Cover the page” questions, so you can check yourself when revising. Answers to the longer questions are in your teacher's answer edition.

LESSON 1: ENERGY STORES AND SYSTEMS

DO NOW

1. For example, a moving car (it is moving) or a hot drink (it is hot). Any sensible answer.
2. joules (J)
3. Warmer: rubbing (friction) transfers energy to the thermal store of your hands.

COVER THE PAGE

1. Joule (J).
2. Any four of: kinetic, gravitational potential, elastic potential, thermal, chemical.
3. A way of transferring energy (by waves).
4. A system that no energy enters or leaves, so its total energy stays the same.

LESSON 2: THE KINETIC STORE

DO NOW

1. The kinetic store.
2. The thermal store (of the brakes and the surroundings).
3. A way energy is transferred (by waves).

COVER THE PAGE

1. Mass and speed.
2. $E_k = \frac{1}{2} m v^2$
3. 0.250 kg
4. It becomes 9 times bigger.

LESSON 3: THE GRAVITATIONAL POTENTIAL STORE

DO NOW

1. Speed, in metres per second (m/s).
2. 0.600 kg
3. $E_k = 0.5 \times 2.0 \times 4.0^2$
 $E_k = 16 \text{ J}$

COVER THE PAGE

1. Gravity pulls each kilogram down with a force of 9.8 N.
2. $E_p = m g h$
3. 98 J
4. $h = \frac{E_p}{m g}$

EQUATION TOOLKIT

Every equation from Lessons 1–3. AQA prints all of them on the equation sheet in the exam, but you still need to know what each symbol means, its unit, and how to use it. Use $g = 9.8 \text{ N/kg}$.

- 1 Cover the quantity you want to find.
- 2 What is left side by side? **Multiply**.
- 3 One above the other? **Divide** the top by the bottom.

Example: to find h from $E_p = m g h$, cover h . E_p is above m and g , so $h = \frac{E_p}{m g}$. Kinetic energy has no triangle: in this unit you only use it to calculate the energy.

$$E_k = \frac{1}{2} m v^2$$

Lesson 2

kinetic energy = $0.5 \times \text{mass} \times (\text{speed})^2$

J, kg, m/s

Use it to calculate the energy. Only the speed is squared.


$$s = v t$$

Lesson 2

distance travelled = speed $\times$ time

m, m/s, s

$$v = \frac{s}{t} \text{ and } t = \frac{s}{v}$$




$$E_p = m g h$$

Lesson 3

gravitational potential energy = mass $\times$ gravitational field strength $\times$ height

J, kg, N/kg, m

$$h = \frac{E_p}{m g} \text{ and } m = \frac{E_p}{g h}$$



1

Write what you know.
Convert first: $g \rightarrow \text{kg}$, $\text{cm} \rightarrow \text{m}$, $\text{kJ} \rightarrow \text{J}$.

2

Write the equation in symbols.

3

Rearrange if you need to, then substitute the numbers.

4

Give the answer with a unit.
Ask: is it sensible?